


Duaklir[®] Pressair[®]

(aclidinium bromide / formoterol fumarate)
inhalation powder 400 mcg/12mcg

A COPD treatment that lets you know when you are using the inhaler correctly.

HOW TO USE THE DUAKLIR PRESSAIR INHALER FOR COPD


1 READY BEFORE INHALATION



Remove the protective cap by lightly squeezing the arrows marked on each side and pulling outwards.



Hold the DUAKLIR PRESSAIR inhaler with the mouthpiece toward you and the orange button facing straight up. Do not tilt.



Press the orange button all the way down and release.



Do not hold the orange button down.
Stop and check to make sure the dose is ready for inhalation.



Check to see if the color of the control window has changed from red to green.



The green control window confirms that your medication is ready for inhalation.

2 CLICK DURING INHALATION



Hold the inhaler way from your mouth, and **breathe out completely.** Never breathe into the inhaler.



Put your lips tightly around the mouthpiece of the DUAKLIR PRESSAIR inhaler. **Take a strong, deep breath.** A “click” sounds during inhalation when the inhaler is used correctly. Keep breathing in, even after you have heard the “click”, to be sure you get the full dose. Remove the DUAKLIR PRESSAIR inhaler from your mouth and hold your breath for as long as it's comfortable. Then slowly breathe out, away from the inhaler.

3 CONFIRM AFTER INHALATION



Make sure the control window has turned from green to red. This “confirms” that you've inhaled your full dose correctly.
If the window did not change to red, you may have forgotten to remove your finger from the orange button before inhaling, or you may not have inhaled correctly. If this happens, go back to Step 2 and repeat the dosing instructions.



Once the control window has turned red, place the protective cap back onto the mouthpiece of the DUAKLIR PRESSAIR inhaler.

INDICATIONS AND USAGE

DUAKLIR PRESSAIR is a prescription medicine to treat chronic obstructive pulmonary disease (COPD). COPD is a long-term (chronic) lung disease that includes chronic bronchitis, emphysema or both.

DUAKLIR is not for the treatment of asthma. DUAKLIR is not a rescue medicine and should not be used for treating sudden breathing problems. Always have a rescue inhaler with you to treat sudden breathing problems. Get emergency medical care if your breathing problems worsen quickly or you use your rescue inhaler, but it does not relieve your breathing problems

IMPORTANT SAFETY INFORMATION

WHO SHOULD NOT USE DUAKLIR PRESSAIR?

Do not use DUAKLIR PRESSAIR if you have asthma without also using a medicine called an inhaled corticosteroid

Do not use DUAKLIR PRESSAIR if you have a severe allergy to milk proteins, or are allergic to aclidinium bromide, formoterol fumarate or any of the ingredients in DUAKLIR PRESSAIR. Ask your healthcare provider if you are not sure.

Please see additional Important Safety Information on the reverse side and accompanying full Prescribing Information.

USEFUL TIPS - Help manage your COPD over the long term



Stick to your treatment plan

Follow your doctor's advice on how and when to take medication. Remember to always use the inhaler correctly for best results



Get regular checkups

Schedule a few visits with your doctor every year -- even if you feel fine! Checking in can help you know if you're on the right treatment plan.



Stay active and eat right

Make sure you stay active and eat right. It can be good for your body. A simple way to start is to take walks and eat a balanced diet.



When you have COPD, quitting smoking is the most important step you can take. If you find it difficult, reach out to a doctor for a professional recommendation.

START SAVING TODAY — THE DUAKLIR \$10 COMMERCIAL GUARANTEE*



If you are an eligible, commercially insured patient, your DUAKLIR PRESSAIR prescription will only cost \$10! Even if DUAKLIR has been denied coverage by your commercial insurance plan.



You may download an active card by visiting Duaklir.com or by asking your healthcare provider.

*Please see Commercial Eligibility rules by visiting Duaklir.com/eligibility-rules/. Restrictions apply.

IMPORTANT SAFETY INFORMATION (CONT'D)

WHAT SHOULD I TELL MY HEALTH CARE PROVIDER BEFORE USING DUAKLIR PRESSAIR?

Tell your health care provider about all your medical conditions, including heart problems, high blood pressure, seizures, thyroid problems, diabetes, eye problems (especially glaucoma), prostate or bladder problems or problems passing urine, are pregnant or plan to become pregnant, or are breast feeding or plan to breast feed.

Tell your health care provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements and eyedrops. DUAKLIR PRESSAIR and certain other medicines may interact with each other and may cause serious side effects. Especially tell your health care provider if you take anticholinergics, other LABA (long-acting beta2-adrenergic agonist) medicines, xanthine derivatives, steroids, diuretics, monoamine oxidase inhibitors, tricyclic antidepressants, beta-blockers or atropine. Use DUAKLIR PRESSAIR exactly as prescribed. Do not use DUAKLIR more often than prescribed or take more medicine than prescribed for you.

WHAT ARE THE POSSIBLE SIDE EFFECT OF DUAKLIR PRESSAIR?

Using too much of a LABA medicine may cause chest pain, increased blood pressure, fast and irregular heartbeat, headache, tremor, or nervousness. Sudden breathing problems may occur immediately after use of DUAKLIR PRESSAIR and may be life threatening. If you have sudden breathing problems immediately after inhaling DUAKLIR PRESSAIR, stop using DUAKLIR PRESSAIR and call your healthcare provider right away or go to the nearest hospital emergency room. Call your doctor or get emergency medical care if you have any symptoms of a serious allergic reaction including rash, swelling of the face and tongue, hives, or breathing problems. You may experience high blood sugar (hyperglycemia) or low blood potassium (hypokalemia). Low blood potassium may cause muscle spasms, muscle weakness, or abnormal heart rhythm. These are not all the possible side effects of DUAKLIR PRESSAIR. Please see full Prescribing Information at Duaklir.com

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Reference: 1. DUAKLIR PRESSAIR (aclidinium bromide/formoterol inhalation powder) [prescribing information]. Morrisville, NC: Circassia Pharmaceuticals Inc;

DUAKLIR PRESSAIR is marketed by Circassia Pharmaceuticals Inc. Circassia is a registered trademark of Circassia Limited. DUAKLIR is a registered trademark of ALMIRALL, S.A. PRESSAIR is a registered trademark of the AstraZeneca group of companies. ©Circassia Pharmaceuticals. All rights reserved. PP-DUA-US-0063 10/19.

